

System of Rice Intensification (SRI) Performance in Morang district during 2005 main season

**Rajendra Uprety, Agriculture Extension Officer
District Agriculture Development Office, Morang, Nepal**

Introduction

Population of the world is still increasing; to feed the larger number of people, the food requirement is increasing day by day, while the land available for food production is decreasing due to losses for new construction and decline in soil fertility. In some areas, the world is changing from chemical-based farming to more organic farming, but still most research and development organizations are packaging and advocating high chemical input-based technology. The prices for chemical fertilizers, seeds of high-yielding varieties, and pesticides are all increasing. This is cutting into farmers' limited purchasing power, and slowly agriculture is becoming a less attractive profession. Many farmers are trying to shift to other businesses. This is the situation of Nepalese farmers and in the rest of the world.

Some decades ago, Nepal was exporting food items to India and other countries. Indeed, rice was the main commodity for export, and there was a government company operating for rice export. Today we are importing food items from India and other countries in high volume. Many of our districts are food-deficit; despite the expenditure of about 200 million rupees, for food transportation and other management costs, we are unable to provide sufficient food to all those districts. Every year we read similar stories of hunger and food deficiency from those districts. In Nepal, rice can be cultivated in most of the districts except Manang and Mustang. Average rice production of Nepal is very low compared with other rice-growing countries around the world. There should be potential to increase rice yields to provide sufficient food for our people and for the export market.

Nepal has 1.56 million hectares of land under rice cultivation, and in 2004, we produced just 4.455 million metric tons rice at the rate of 2.86 tons/ha. Out of this area, 1 million hectares have irrigation facilities. The District Agricultural Development Office (DADO), Morang and some other organizations have been trying out the System of Rice Intensification (SRI) method for rice cultivation in Nepal. This method has been showing remarkable potential to reduce hunger and food-deficiency problems in Nepal. This report gives some results of main season rice 2005 of Morang and a few other districts.

Morang is the largest rice producer district of Nepal locates in the eastern terai. It cultivates rice in more than 94 thousand hectares and produces more than 311 thousand metric tons of rice. Nearly half of its lands have permanent irrigation facility. Its average production during last some years has slightly more than 3 metric tons/hectar. All rice growing village development committees are located in terai. DADO Morang has been trying SRI method for last three years. Our focus is more demonstrative for dissemination this method than research.

In Morang district, we started SRI three years ago on a small plot (just about 100 square meters) using a handful of rice seeds (Radha 12 variety). The result of that plot encouraged us to disseminate this method to other parts of the district. In the beginning, our pace has slow, but

after we got support from CIIFAD, we could speed up our activities and expanded the areas of SRI cultivation. Last year, SRI results made DADO/Morang a successful entrant in the 2005 Nepal Development Marketplace competition, organized jointly by the Kathmandu office of the World Bank and the Poverty Alleviation Fund. Out of 1137 applicants, our SRI Promotion Project was one of 20 selected for funding.

As Project team leader, I have the challenge and opportunity now to prove the potentials and suitability of SRI methods for our climate and conditions. I have modest funds for project activity and some experienced extension workers and innovative leader farmers to work with. The organization and management of these resources (human and capital) have achieved a good result in first season of our project.

During our first season with NDM support, we hosted various visitors from the east (Mechi) to the west (Mahakali) of our country. Among the visitors of our SRI farmers' fields were the Hon. Minister of Agriculture and Cooperatives (Mr. Badri Prasad Mandal), Joint Secretaries of the Ministry of Agriculture and Cooperatives, Directors-General of the Departments of Agriculture and Livestock Services, Executive Director and Directors of the Nepal Agriculture Research Council (NARC), Executive Director and board members from the Poverty Alleviation Fund, visitors from the World Bank/Kathmandu office, and several other seniors officers, technicians and farmers from different organizations and districts. Besides these visitors, we also received renowned journalists such as Mr. Kunda Dixit (editor, *Nepali Times*), Mr. Rajendra Dahal (editor, *Himal Khabarpatrika*), Mr. Charles Haviland (Nepal correspondent, BBC World Service) and several others from NTV, Kantipur, Gorkhapatra, Channel Nepal, etc.

Methodology

Our activities have been focus on demonstration and dissemination of SRI method in larger areas. In same time we have been collecting maximum information related with this method. Most of the new farmers start SRI in small area (like 1-2 Kathas) and after gain some confidence increased the areas under this method. Information has collected by used DADO extension workers and leader farmers through designed format from the field.

We recommend three spacing for transplanting 30x30 cm., 25x25 cm. and 20x20 cm. according to crop duration and tillering habits of the variety. But some farmers also used 25x20 cm spacing. Most of the farmers used seedling within 12 days but some farmers also used older seedlings due to delay monsoon. For water management we used flooding and drying system, in case of clay soil we recommended to maintain moisture level soil (not stagnant of water) without drying the during vegetative stage. For loose, high organic matter contained loam soil has dried up to cracking level 4-5 times. Fertilizers and manure dose has open for farmers as they were using with traditional method. We have found 15 varieties of rice have been used with SRI method. Main varieties information has given in this report.

For yield record, total crop harvest for small plots and 5 (2x2.5 meters) square meters areas yield was taken for larger plots (from 2-3 places). For moisture adjustment, we reduce 13-14 % moisture from the recorded yield to finalize the yield data (moisture testing by used moisture meter we found, in general 27-28 % moisture in SRI rice).

Objectives

Basic objective of our activities is to increase rice yield in our present inputs level of our small and marginal farmers, who cultivate their land themselves. Besides this we have some other objectives of our program, as

1. To disseminate SRI method in larger area of Morang and other parts of the country.
2. Exploring maximum information related with SRI method, and packaging appropriate package of practice for different varieties and soil condition.
3. To bringing SRI method in mainstream of agriculture development endeavors.

Results and Discussion

In the previous year, 2004, we had about 2 dozen farmers in 16 Village Development Committee (VDC) areas; in the early season of this year, 2005, we have about 6 dozen farmers in 22 VDCs. Then in the main season 2005, with financial support from the NDM, we had more than than 1400 farmers in 53 VDCs of Morang district, and about 100 farmers in 9 VDCs of neighboring Panchthar district participating. To evaluate performance of SRI methods, we collected data from >400 SRI farmers, and this information is presented in this report.

Table 1. Area under SRI method in Morang district (2005)

SN	Category	Number	Percentage
1	<2 Kathas	35	8.5
2	2 Kathas	287	69.5
3	2.1-5 Kathas	70	16.8
4	>5 Kathas	21	5.2
Range: 1-65 Kathas			
Average: 2.84 Kathas		413	100

System of Rice Intensification (SRI) method is still new for Nepal and Morang farmers. But the numbers and areas of SRI farming is increasing season after season, going in 5 seasons (two and a half years) from just 1 farmer using the methods, to 1400 farmers in 53 VDCs, with farm size under SRI method ranging from 1 Katha (333 square meters) to 65 Kathas (2.16 hectares). Generally new farmers have been trying SRI first on a small scale, and after they gained experience they increased their area under SRI method.

Table 2. Rice varieties used with SRI method, Morang district (2005)

SN	Rice Variety	Number	Percentage	Average Production by SRI (Mt./ha)
----	--------------	--------	------------	------------------------------------

1	<i>Bansdhan</i>	248	59.8	6.36 (2.5-11.0)
2	<i>Mansuli</i>	48	11.6	5.74 (3.6-9.9)
3	<i>Suwarna</i>	40	9.6	6.48 (3.5-11.0)
4	<i>Hardinath 1</i>	39	9.4	5.82 (3.6-8.4)
5	<i>Barse 2014</i>	13	3.1	5.5 (4.2-6.8)
6	<i>Suganda</i>	12	2.9	5.05 (3.0-7.5)
7	<i>Radha 12</i>	12	2.9	7.44 (5.2-10.0)
8	<i>Pusa 834</i>	1	0.2	6.0
9	<i>Barse 3017</i>	1	0.2	7.1
10	<i>Basmati</i>	1	0.2	3.9 (FP = 2.4)
		415	100	Average: 6.3 Mt/ha

Several varieties of rice have been used with SRI method. Among them *Bansdhan* or *Kanchi mansuli* was by far the most popular one in Morang district. SRI users' preference was also for *Mansuli*, *Suwarna*, and *Hardinath* (arecently released 120-day variety). Regarding average production, *Radha-12* ranked first, followed by *Suwarna* and *Bansdhan*, respectively. But *Suwarna* and *Bansdhan* had the highest recorded yields, up to 11 Mt/ha. That *Radha-12* had the highest average yield was due to a lesser number of farmers using this variety.

Among varieties with fine quality, scented rice, *Suganda*, *Basmati* and *Pusa-835* also produced good yield with SRI compared with farmers' practice. This has shown possibility of profitability with lower-yielding but high-quality and high-price rice. Generally such varieties have lodging problems under usual management practices. But with SRI methods, these lodging problems are solved, and single seedlings have produced 21.8 panicles/hill (maximum 46 and minimum 13) with an average of 159 grains/panicle, the highest average number of grains/panicle (213) in black *Basmati* rice.

Table 3. Age of seedlings used with SRI method, Morang district (2005)

SN	Seedlings age (days)	Number	Percentage	Average Production by SRI method (Mt/ha)
1	8	22	5.4	6.94 (4.0-9.0)
2	9-10	123	30.0	6.32 (3.6-11.0)
3	11-12	169	41.3	6.41 (2.5-11.0)
4	13-14	64	15.6	5.77 (3.0-9.0)
5	15 and above	32	7.8	5.52 (3.7-7.0)
Range: 8-21 days				Average: 6.3 Mt/ha
Average: 11.4 days		410	100	

SRI method generally recommends 8-12 day-old seedlings. But this year, due to a late monsoon and lack of irrigation facilities, some of our farmers have planted older seedlings while using the other SRI practices (single seedling, spaced planting, active weeding, with less water). Still, the majority of SRI farmers have transplanted their crop within the age of 12 days. Highest average production has harvested from the youngest aged seedlings, and lowest average was from older-aged seedlings. But highest individual yields were achieved from the 9-12-day group, due to damage of some younger seedlings from the heavy rain during transplanting time and farmers didn't fill the gaps later on. From our experience, we see that if we undertake timely

weeding and do good water management, we can produce improved yield from seedlings up to 15 days old with SRI methods. Besides seedling age, rice yield has influenced by weed management, organic manure, water management, soil structure, and soil fertility.

Table 4. Number of times of weeding by SRI farmers, Morang district (2005)

SN	Times of weeding	Number	Percentage	Productivity of SRI method Mt/ha
1	One	32	8.0	5.16 (range 3.6-7.6)
2	Two	366	88.6	5.87 (range 3.5-11)
3	Three	14	3.4	7.87 (range 5.85-10.4)
Average: 1.9 times		412	100	Average: 6.3 Mt/ha

Weeding is a very important practice for SRI method. It directly influences rice yield and earlier maturity. The average production of rice by those who used SRI methods was highest with three weeding, followed two and single weeding. The highest yield 11 Mt/ha was found with 2 weeding, due to other factors like soil quality and other management practice. That an additional weeding, going from two to three, added an average of 2 t/ha to yield makes this additional investment of labor very profitable.

In general we found less weed problems in soil with high organic matter content. But weed control methods with the rotating hoe give benefits most of the time with SRI, because of its effect on soil aeration. Early weeding enhances production of more primary tillers, which ultimately produces larger panicles having more grains and higher yield. If weeding is begun late, then the number of weeding may not be so beneficial. If we keep our crop weed-free for the first month after transplant, weeding will have a positive influence on rice yield. *What is not measured and evaluated in these data is the effect that soil-aerating weeding has on soil biota (organisms) which provide numerous services to the growing rice plants. This is an area for future investigation.*

Table 5. Yield distribution by SRI method, Morang district (2005)

SN	Yield of SRI method (Mt/ha)	Number	Percentage
1	Below 4	21	5.1
2	4.1-5	69	16.7
3	5.1-6	107	26.0
4	6.1-7	112	27.3
5	7.1-8	64	15.5
6	8.1-9	26	6.3
7	9.1-10	9	2.2
8	10.1 and above	4	0.9
Range: 2.5-11.00 Mt/ha		412	100
Average: 6.3 Mt/ha			

With SRI methods, farmers in Morang district using them produced about double as much as much rice as with usual methods in the 2005 main season. Previously these farmers produced 3.1 metric tons of rice on average with conventional methods. The range of SRI productions was 2.5 metric tons to 11.0 metric tons, and average production was 6.3 metric tons

by SRI methods. The majority of SRI farmers produced more than 6 metric tons yield, and very few got less than 4 metric tons.

Table 6. Yield distribution by farmers' method, Morang district (2005)

SN	Yield of farmer's method (Mt/ha)	Number	Percentage
1	Up to 3	258	62.6
2	3.1-4	131	31.8
3	4.1-5	21	5.1
4	5.1 and above	2	0.5
Range: 1.8-6.0 Mt/ha Average: 3.1 Mt/ha		412	100

The range of production achieved with farmer's usual methods was 1.8 to 6.0 metric tons (Table 6). But a majority of them (62.6 %) produced under 3 metric tons and less than one percent of farmers (0.5 %) produced more than 5 metric tons by using farmers' method.

Table 7. Yield difference between SRI and farmers' methods, Morang district (2005)

SN	% Yield difference in favor of SRI	Number	Percentage
1	Up to 50%	27	6.6
2	51-75%	63	15.3
3	76-100%	142	34.5
4	101-125%	96	23.3
5	126-150%	64	15.5
6	151-175%	10	2.4
7	> 175%	10	2.4
Range: 11.1-205.6% Average: 99.8%		412	100

Range of yield difference between SRI and farmers has been found very large. The yield difference between these two methods was from 11 to 205 percent, and the average was 99.8%. Only 6.6 % farmers had less than 50 % difference, and almost 5 % farmers produced more than 150 % more by using SRI method. Table 7 gives details.

Table 8. Age of seedlings and crop duration (var. *Bansdhan*) with SRI method, Morang district (2005)

SN	Seedling age (days)	Farmers Number	Average crop duration (days)	Crop duration with usual methods (days)	Earliness in maturity with SRI (days)
1	8-9	5	123.6	145	21.4
2	10-11	14	129.7	145	15.3
3	12-14	23	131.6	145	13.4
4	15 and above	9	138.5	145	6.5
		51			

Bansdhan is a most popular and most widely planted variety of rice in Morang and the eastern terai of Nepal. This year we analyzed earliness in maturity of rice with respect to seedling age. According to the data in Table 8, we see a definite positive relationship between seedling age and earliness when SRI practices are used. Younger seedlings mature earlier; it may

be due to early initiation and completion of vegetative stage and entering into the reproductive stage. This can be verified by conducting systematic research on this in the coming season.

Table 9. Age of seedling and yield of rice by SRI method, Morang district (2005)

S N.	Age of Seedling	Yield of rice (Mt/ha)							
		Up to 4	4.1-5	5.1-6	6.1-7	7.1-8	8.1-9	9.1-10	>10
1	8	1	1	2	6	5	6	0	0
2	9-10	5	21	34	29	25	4	5	2
3	11-12	11	18	40	54	29	12	4	2
4	13-14	2	16	22	17	2	4	0	0
5	15 & above	2	12	6	6	3	0	0	0
Total		21	68	104	112	64	26	9	4

Age of seedling is a very important component of the SRI methodology. In this study, the effect of age of seedling on the yield of rice was analyzed. The above table shows the relationship between age of seedling and yield of rice produced by use of SRI methods. A majority of farmers who used younger seedlings produced more than 6 metric tons per hectare (81.9% of those using 8-day-old seedlings and 52.8% of those with 9-10-day-old seedlings). On the other hand, a majority of those farmers who used older seedlings produced less than 6 tons per hectare. The data in the above table indicate that seedling age between 8-12 days is best for SRI methods. Older seedlings up to 15 days can produce more yields compared to traditional methods, but their yield improvement is generally less.

Table 10. Number of weedings and yield of rice by SRI method, Morang district (2005)

S N.	Numbers of weeding	Yield of rice (Mt/ha)							
		Up to 4	4.1-5	5.1-6	6.1-7	7.1-8	8.1-9	9.1-10	>10
1	1	6	5	11	8	2	0	0	0
2	2	15	64	95	100	57	26	6	3
3	3	0	0	1	4	5	0	3	1
Total		21	68	107	112	64	26	9	3

Weeding is another important operation to increase yields with SRI methods. The relationship between number of weedings and rice yield is shown in Table 10. With a single weeding, the majority of SRI farmers (68.75%) produced at least 6 tons of yield per hectare. But with two weedings, a majority of farmers (52.2%) produced more than 6 tons yield. Even more impressive, with three weedings, most of the farmers (92.8%) produced more than 6 tons yield, and 28.6% farmers produced more than 9 tons yield per hectare. Next season, we will encourage some farmers to do four weedings if possible, to see what further gains in yield might be possible as a result of active soil aeration.

Table 11. Age of seedling and early maturity by SRI method, Morang district (2005)

S N	Age of Seedling (days)	Earliness in maturity (days)					
		Up to 5	6-10	11-15	16-20	21-25	>25
1	8	0	2	2	10	8	0
2	9-10	1	16	36	31	27	12
3	11-12	5	21	44	49	38	12

4	13-14	1	9	17	23	11	3
5	15 and above	11	6	7	5	3	0
Total		18	54	106	118	87	27

Short crop duration is one benefit we can find from using SRI methods. Most varieties will mature earlier this way than when grown with conventional methods. A majority of the SRI rice crop (81.8%) with 8-day-old seedlings matured 16 days and more earlier, and >36% of the crop matured more than 20 days earlier compared with traditional methods. Similarly, 56.9% of SRI crops with 9-10-day-old seedlings matured more than 16 days earlier. But, in the case of older aged seedlings (15 days and above), only 25% of the crops are matured 16 or more days earlier, while but 34.3% of such SRI crops were earlier by up to 5 days. Details are presented in Table 11. It is a real benefit for farmers to be able to harvest their crop earlier, reducing risks of loss or damage and early planting of the next crops.

Table 12. Time of first weeding and early maturity by SRI method, Morang district (2005)

S N	Time of first weeding (days)	Earliness in maturity (days)					
		= 5	6-10	11-15	16-20	21-25	26 & more
1	Up to 10 th day	0 (0%)	3 (23%)	3 (23%)	4 (31%)	2 (13%)	1 (8%)
2	11 th -15 th day	0 (0%)	8 (18%)	11 (25%)	15 (33%)	9 (20%)	2 (4%)
3	16 th -20 th day	2 (2%)	10 (10%)	23 (23%)	29 (29%)	30 (30%)	5 (5%)
4	21 st -25 th day	6 (7%)	13 (16%)	24 (30%)	19 (24%)	10 (13%)	8 (10%)
5	26 th day or later	8 (12%)	5 (8%)	18 (27%)	17 (25%)	17 (25%)	2 (3%)
Total		16	39	79	84	68	18

Early start of weeding has been useful for promoting early maturity and more production. In Table 12, the relationship between date of first weeding and early maturity is presented. Data indicate that early weeding enhances earlier maturation of the crop. Early weeding helps stimulate early tillering, and early tillering helps the early vegetative stage and ultimately earlier maturity in SRI crops. This can be verified through systematic research in the coming season.

Table 13. Number of weedings and early maturity by SRI method, Morang district (2005)

S N	Numbers of weeding	Earliness in maturity (days)					
		= 5	6-10	11-15	16-20	21-25	26 & more
1	One	1 (3%)	6 (18%)	11 (34%)	8 (24%)	6 (18%)	0
2	Two	17 (5%)	48 (13%)	93 (26%)	104 (28%)	77 (21%)	25 (7%)
3	Three	0	0	2 (14%)	6 (43%)	4 (29%)	2 (14 %)
		18	54	106	118	87	27

Getting more rice production from more weeding is often reported with SRI crops around the world. When using a rotary hoe, this does not only control weeds but it aerates the top horizon of the soil, which stimulates the growth of aerobic bacteria and fungi in the soil. In Table 13, we relate the number of weedings with earliness of maturity for rice crops. The data show a definite relationship between the number of weedings and earliness/short crop duration in rice crop.

With a single weeding, a difference in crop duration between SRI and traditional farming methods is observable but not so great as when the number of weedings is increased. This could become one of the most cost-effective components of SRI practice, since normally farmers have been applying expensive fertilizer to raise yield, when with some labor and skill they could get as large or larger increments by stimulating the services of soil biota, usually suppressed under the anaerobic conditions of continuous flooding which is farmer practice.

Table 14. Seed rate used by SRI farmers in Morang district (2005)

SN	Seed rate (Kg./ha)	Number	Percentage
1	Up to 3	35	8.6
2	3.1-4	96	23.59
3	4.1-5	116	28.53
4	5.1-6	108	26.53
5	6.1-7	5	1.2
6	>7.1	47	11.55
Range: 2.3-7.5 Kg./ha.			
Average: 4.9 Kg/ha.		407	100
Seed rate used by farmers: 60-75 Kg./ha.			

By used SRI method farmers can reduced seed rate drastically. In Morang, average seed rate used by SRI farmers was 4.9 Kg./ha. (Only 8.16 %) and lowest seed rate was found 2.3 Kg./ha (3.83 %) and highest seed rate was 7.5 Kg./ha (Only 12.5 %). In this way farmers can save more than 90 percent seed for their consumption or other use.

Table 15. Fertile tillers and grains of different rice varieties in SRI method, Morang district (2005)

SN	Rice Variety	Average Fertile tiller/hill			Average numbers of grain/panicle		
		Average	Maximum	Minimum	Average	Maximum	Minimum
1	<i>Bansdhan</i>	19.72	47	7	195	404	62
2	<i>Mansuli</i>	19.62	48	7	195.5	390	50
3	<i>Suwarna</i>	17.5	30	9	197	352	60
4	<i>Hardinath 1</i>	11.9	16	7	168.6	257	51
5	<i>Barse 2014</i>	20	37	9	179.5	286	57
6	<i>Suganda</i>	12.5	17	7	159	230	105
7	<i>Radha 12</i>	18.7	33	10	211.7	345	68
8	<i>Pusa 834</i>	17	22	12	149	200	52
9	<i>Barse 3017</i>	28	-	-	199	271	96
10	<i>Basmati</i>	21.8	46	11	158.5	213	102
		18.77	29.9	(12.7)	192.5	291	(87)

SRI method has found very helpful to increase fertile tiller numbers, numbers of grain per panicle. Table 14 is presenting details of fertile tiller (panicle numbers/hill) and numbers of grain/panicle. In SRI method we have found larger panicles, more numbers of grains and larger grains. In observation we found that early tillering produce larger panicle and more grain numbers compare to secondary and tertiary tillers. To encourage more early tillers, early

weeding has found very useful. To verify relationship between weeding and panicles we need to conduct systematic research in coming season.

Conclusion

Three years results and experiences has shown that SRI method is very useful for small and marginal framers, who cultivate their land themselves. It has reduced use of seed, water, and pesticides for more production. By used locally available seed and other inputs farmers can produce about double production from same land. With better management farmers can triple their production. Besides these benefits farmers can harvest their rice crop 15-25 days earlier, which will help in early planting of the next crop (as wheat, maize, oil seed, and vegetables) and enhance to produce more yield from those crops. Weed management has found the main constraints for SRI method. About 20-30 labors (man days) more, necessary for SRI method compare to traditional method. But it will be reduced next season. If we properly manage weeds this year, weed population in next season automatically reduced (due to less weed seed for germination). Besides this some farmers used chemical control with one manual weeding to reduce labor cost. Use of mechanical weeder will be another remedies for the reduction of labor cost.

Recommendation for SRI users

Based on our experience we can recommend some practice of package for farmers' application.

1. Age of seedling should be within 10-12 days for better yield and early harvesting. But we can use up to 15 days seedlings successfully in this method.
2. Dry seedbed is always useful for seedling rising, solarization technique will be very helpful for early, vigor and healthy seedling.
3. Spacing of plant should be maintain according to varieties (duration, tillering habit), soil and water management, same spacing for all varieties and situation will not be useful.
4. In case of clay soil some moisture should be maintain during vegetative stage, in high organic matter contained loam soil we can use wet and dry practice (drying up to cracking level) for better result.
5. Weed management in early stage of vegetative growth has found very useful for more fertile tiller numbers as well as for more and larger grains. Farmers should keep their SRI field weed free for first 4-6 week after transplanting.
6. Organic matter/manures has found very useful for SRI method but in case of less fertile soil farmers should use balance dose of fertilizers for more yield.

Recommendation for future research

Some vary interesting findings has found during our observation and data has shown such relationship between practice of SRI and yield and earliness of rice crop. To verify these relationships we need to conduct some research in coming season. Some recommendations are as follows:

1. Relationship between seedling ages with yield and earliness of crop maturity.
2. Relationship between time and numbers of weeding with yield and earliness of crop maturity.
3. Relationship between organic manure and fertilizers with yield and earliness of crop maturity.
4. Relationship of water management and yield of SRI method.

System of Rice Intensification (SRI) Method in Panchthar (2005 Main Season)

Total Farmers: 95

VDCs: 9

Varieties used with SRI: 12 (Ramtulabi, Masino, Jamara, Phulpata, Khumal-4, Sabitri, Arun, Chirankhe, Tauli, Champe, Bhangari, Chinapate).

Fertilizers Used :0

Compost: 0 (used previous crop Maize).

Seed rate: 5-8 Kg/Ha (average-5.99 Kg/ha).

Spacing: 30x25 cm., 25x25 cm.

Weeding: 2-3 times.

Seedling age: 11.7 days (8-18 days)

Average Panicle/hill: 16.4 (13-32).

Average Grains/panicle: 209.4

Average Production with SRI method: 4.6 Mt./ha. (3.74-6.48 Mt./ha)

District Average Production: 2.1 Mt./ha.

****Source: DADO Panchthar.**

System of Rice Intensification (SRI) Method in Sunsari (2005 Main Season)

VDC: 2 (Chimdi, Amahibelaha)

Farmers: 9

Varieties Used: 2 (Sugandha-1, Hardinath-1)

Average Production: 5.6 Mt./ha (4.3-6.7 Mt./ha)

Average production of that area: 2-3 Mt/ha

Average of Sunsari district: 3.12 Mt./ha

****Source: LI-BIRD, Site office Sunsari**